Subduing Photosensitive Dermatitis through Ayurveda- Clinical images

Rashmi TM^{1*}, Sathish HS² ¹ Assistant Professor, Dept. of *Kayachikitsa*, ² Associate Professor, Dept. of *Shalya Tantra*, Indian Institute of *Ayurveda* Research and Hospital, Rajkot, Gujarat, India *Corresponding author: email: ayurrashmitm@gmail.com Mob: 09624459009

Abstract:

No abstract is required as this is clinical images.

Key words: Ayurveda, Dermatitis, Kushta, Panchakama, Shodhana, Virechan

A male patient aged 54 yrs, working as a farmer approached with severe itching sensation all over the face and neck with thick blackish discolouration since 2 yrs and with futile results with the previously taken medications [Tab Cetrizine 5 mg]. Itching is so severe that the patient always used to keep a thick napkin with him and keeps rubbing all over the face and neck only to get a little solace. History revealed that he is suffering with the above complaints since 2 years, which was less severe and only on forehead initially and later manifested all over the face and neck with intense itching. No history of any other systemic diseases noted. On local examination skin looks thick, cracked with dark discolouration. Continuous rubbing due to severe itching has led to local inflammation also which was pronounce on eyelids (Fig-1). Case was diagnosed as Tridoshaja Kushta with the predominance of Vata and Kapha and as photosensitive dermatitis from contemporary perspective.

Plan of treatment:

- ✓ Virechana followed by Shamanoushadhi: Deepana pachana was done with Amapachana Vati 500 mg 1 TID for 3 days.
- ✓ Started *Snehapana* with *Guggulu Tikthaka Ghritha*, 30 ml on first day.
- ✓ Snehapana was given up to 6 days with 200 ml as maximum dose of sneha on 6th day.
- ✓ Sarvanga abhyanga with Nimba Taila and Sarvanga Bashpa sweda done for 3 days.
- ✓ Subsequent day, Sarvanga Abhyanaga followed by Bashpa Sweda was administered and orally Trivrut Lehya 50 gms was administered for Virechana purpose.
- ✓ 21 vegas [Frequency of Purgation] were recorded, no untoward events recorded during Shodhana.
- ✓ Proper Samsarjana krama (Post virechan measures) was advised and patient discharged with the following Shamanoushadhi (internal medication):
 - Arogyavardhini Vati 250mg 2 TID [Before food with plain water].

INTERNATIONAL JOURNAL OF AYUSH CASE REPORTS website: www.ijacare.in

ISSN: 2457-0443

- Sarivadi Vati 250mg 2 TID [After food with warm water].
- Mahatiktha Ghritha 3tsp BID [After food with warm water].
- Treatment Period 2 months. \checkmark

Observations:

During Shodhana itching was reduced almost up to 70% on day second of

Clinical Images:

After shodhana snehapana only. inflammation was completely reduced especially on eyelids (Fig-2). After 2 months treatment there was no itching at all. Thickness of skin and blackish discolouration also reduced significantly (Fig-3).




Fig. 2. After 1 month of treatment

INTERNATIONAL JOURNAL OF AYUSH CASE REPORTS website: www.ijacare.in ISSN: 2457-0443



Guarantor: Corresponding author is guarantor of this article and its contents.

Conflict of interest: Author declare that there is no conflict of interest

How to cite this article:

Rashmi TM, Sathish HS. Subduing Photosensitive Dermatitis through Ayurveda - Clinical images. Int. J AYUSH CaRe. 2017;1(2): 37-39